## Table 24.1 Cancer of the Stomach (Invasive)

Trends in SEER Incidence<sup>ab</sup> and U.S. Mortality<sup>c</sup> Using the Joinpoint Regression Program, 1975-2008 With up to Five Joinpoints, 1992-2008 With up to Three Joinpoints, Both Sexes by Race/Ethnicity

	JP Trend 1	JP Tr	JP Trend 2		end 3	JP Trend 4		JP Trend 5		JP Trend 6		AAPC <sup>d</sup>	
	Years APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	1999-08	2004-08
SEER 9 Delay-Adju	sted Incidencea, 1	975-2008											
All Races	1975-08 -1.6*											-1.6*	-1.6*
White	1975-08 -1.8*											-1.8*	-1.8*
Black	1975-08 -1.3*											-1.3*	-1.3*
SEER 13 Delay-Adj	usted Incidenceb,	1992-2008											
All Races	1992-08 -1.3*											-1.3*	-1.3*
White	1992-08 -1.3*											-1.3*	-1.3*
Black	1992-08 -1.6*											-1.6*	-1.6*
SEER 9 Observed Incidence <sup>a</sup> , 1975-2008													
All Races	1975-08 -1.6*											-1.6*	-1.6*
White	1975-08 -1.8*											-1.8*	-1.8*
Black	1975-08 -1.3*											-1.3*	-1.3*
SEER 13 Observed Incidence <sup>b</sup> , 1992-2008													
All Races	1992-08 -1.4*											-1.4*	-1.4*
White	1992-08 -1.4*											-1.4*	-1.4*
White NH <sup>ef</sup>	1992-08 -1.8*											-1.8*	-1.8*
Black	1992-08 -1.7*											-1.7*	-1.7*
Black NH <sup>ef</sup>	1992-08 -1.7*											-1.7*	-1.7*
API <sup>e</sup>	1992-08 -3.0*											-3.0*	-3.0*
AI/AN <sup>eg</sup>	1992-08 -0.4											-0.4	-0.4
Hispanic <sup>f</sup>	1992-08 -1.5*											-1.5*	-1.5*
U.S. Cancer Morta	lity <sup>c</sup> , 1975-2008												
All Races	1975-87 -2.6*	1987-90	-0.2	1990-08	-3.0*							-3.0*	-3.0*
White	1975-87 -2.8*	1987-90	-0.7	1990-08	-3.2*							-3.2*	-3.2*
Black	1975-87 -1.6*	1987-91	0.6	1991-08	-3.0*							-3.0*	-3.0*
U.S. Cancer Morta	lity <sup>c</sup> , 1992-2008												
All Races	1992-08 -3.1*											-3.1*	-3.1*
White	1992-08 -3.2*											-3.2*	-3.2*
White NH <sup>ef</sup>	1992-94 -1.4	1994-08	-3.6*									-3.6*	-3.6*
Black	1992-99 -2.2*	1999-06	-3.9*	2006-08	-0.4							-3.1*	-2.2
Black NH <sup>ef</sup>	1992-08 -2.8*											-2.8*	-2.8*
API <sup>e</sup>	1992-08 -3.7*											-3.7*	-3.7*
AI/AN <sup>eg</sup>	1992-02 0.9	2002-08	-7.0*									-4.5*	-7.0*
Hispanic <sup>f</sup>	1992-04 -1.9*	2004-08	-5.2*									-3.4*	-5.2*
-													

Joinpoint Regression Program Version 3.5, April 2011, National Cancer Institute. (<a href="http://surveillance.cancer.gov/joinpoint/">http://surveillance.cancer.gov/joinpoint/</a>).
The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The Hispanic and Non-Hispanic mortality trends exclude deaths from CT, DC, ME, MD, MN, NH, NY, ND, OK and VT.

Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

The APC/AAPC is significantly different from zero (p<.05).

<sup>-</sup> Joinpoint regression line analysis could not be performed on data series.